

Supplementary Material

10.1302/0301-620X.104B1.BJJ-2021-0325.R2

Table i. Radiostereometric migration results from studies on force-closed design.

Stems with force-closed design	Subsidence at 24 mths (mm)	Retroversion at 24 mths (°)
CPS-plus* (Glyn-Jones et al) ¹	0.67	0.03
CPCSt (McCalden et al) ²	0.77	1.61
C-stem‡ (Sundberg et al) ³	1.35	1.9
C-stem (von Schewelov et al) ⁴	1.36	1.6
Exeter§ (Stefansdottir et al) ⁵	1.34	1.2
Exeter§ (McCladen et al) ²	1.25	0.59
Exeter§ (Alfaro-Adrian et al) ⁶	1.2	1.12
Exeter§ (Glyn-Jones et al) ¹	0.87	1.28
Exeter§ (Murray et al) ⁷	0.7	0.7
Corail line to line¶ (present study)	0.86	1.13
Corail standard¶ (present study)	1.22	2.36

*Smith & Nephew Orthopaedics, Switzerland.

† Smith & Nephew, USA.

‡ DePuy, USA.

§Stryker, USA.

¶DePuy, Warsaw, Indiana, USA.

References

1. Glyn-Jones S, Gill H, Beard D, McLardy-Smith P, Murray D. Influence of stem geometry on the stability of polished tapered cemented femoral stems. *J Bone Joint Surgery Br.* 87(7):921–927.
2. McCalden R, Charron K, Yuan X, Bourne R, Naudie D, MacDonald S. Randomised controlled trial comparing early migration of two collarless polished cemented stems using radiostereometric analysis. *J Bone Joint Surgery Br.* 92(7):935–940.
3. Sundberg M, Besjakov J, von Schewelov T, Carlsson A. Movement patterns of the c-stem femoral component: An RSA study of 33 primary total hip arthroplasties followed for two years. *J Bone Joint Surgery Br.* 87-B(10):1352– 1356.
4. von Schewelov T, Carlsson A, Sanzén L, Besjakov J. Continuous distal migration and internal rotation of the c-stem prosthesis without any adverse clinical effects: An RSA study of 33 primary total hip arthroplasties followed for up to ten years. *Bone Joint J.* 2014;96-B(5):604–608.
5. Stefánsdóttir A, Franzén H, Johnsson R, Ornstein E, Sundberg M. Movement pattern of the Exeter femoral stem; a radiostereometric analysis of 22 primary hip arthroplasties followed for 5 years. *Acta Orthop Scand.* 2004;75(4):408–414.
6. Alfaro-Adrian J, Gill H, Murray D. Cement migration after THR: a comparison of Charnley Elite and Exeter femoral stems using RSA. *J Bone Joint Surgery Br.* 81(1):130–134.
7. Murray DW, Gulati A, Gill HS. Ten-year RSA-measured migration of the Exeter femoral stem. *Bone Joint J.* 2013;95-B(5):605–608.