

# Bone & Joint Open



**Supplementary Material**

**Table i.** Comparison of patient-reported scores of those waiting for total joint replacement for those taking opioid and those not taking opioids.

Variable	Taking opioid	No opioid	Total	p-value
Median EQ-5D Index; IQR (range)	-0.038; 0.20 to 0.21 (0.59 to 0.74)	0.206; 0.10 to 0.41 (0.59 to 1.00)	0.155; -0.13 to 0.37 (-0.59 to 1.00)	< 0.001**
WTD state (n, %)	197 (35.5)	80 (51.6)	277 (39.0)	< 0.001##
Median EQ-5D VAS; IQR (range)	35.0; 20.0 to 60.0 (0 to 100)	45.0; 25.0 to 60.0 (0 to 100)	40.0; 20.0 to 60.0 (0 to 100)	0.018**
Median Hip Oxford Total Score; IQR (range)	7.0; 4.0 to 12.0 (0 to 37)	10.0; 5.0 to 18.0 (0 to 46)	9.0; 5.0 to 18.0 (0 to 46)	0.038**
Median Hip Oxford Pain Score; IQR (range)	5.0; 5.0 to 5.0 (3 to 5)	5.0; 4.0 to 5.0 (1 to 5)	5.0; 4.0 to 5.0 (1 to 5)	0.024**
Median Knee Oxford Total Score; IQR (range)	7.0; 5.0 to 13.0 (0 to 33)	11.0; 6.0 to 18.0 (0 to 48)	10.0; 5.0 to 17.0 (0 to 48)	< 0.001**
Median Knee Oxford Pain Score; IQR (range)	5.0; 5.0 to 5.0 (1 to 5)	5.0; 4.0 to 5.0 (1 to 5)	5.0; 4.0 to 5.0 (1 to 5)	0.007**

\*Significant difference between those taking opioids and those not taking opioids.

#Chi-squared test.

+Mann-Whitney U Test

EQ-5D, EuroQol five dimension; IQR, interquartile range; VAS, visual analogue scale.