

We welcome letters to the Editor concerning articles which have recently been published. Such letters will be subject to the usual stages of selection and editing; where appropriate the authors of the original article will be offered the opportunity to reply.

Letters should normally be under 300 words in length, double-spaced throughout, signed by all authors and fully referenced. The edited version will be returned for approval before publication.

©2005 British Editorial Society of Bone and Joint Surgery
J Bone Joint Surg [Br] 2005;87-B:884-6.

Predictive value of the duration of sciatica for lumbar discectomy

Sir,

I read with interest the article by Ng and Sell¹ in the May 2004 issue entitled "Predictive value of the duration of sciatica for lumbar discectomy". It is a rigorous assessment of this much debated question, but to call it a prospective, cohort study, despite the fact that the duration of symptoms varies at recruitment, may lead readers to draw a possibly erroneous conclusion.

It is the nature of sciatica from disc prolapse that patients may improve at any time, with a gradual decrease in the incidence of patients experiencing sufficient improvement to obviate the need for surgery. The necessary corollary of this is that the longer a patient has had sciatica, the nearer is he or she to the worse end of the spectrum of disease. This may be the entire explanation of the worse outcome - by waiting longer, one is not predicting a worse result from surgery, but merely selecting a worse group of patients.

What is needed is a cohort study with recruitment based on an intention to treat (by any means - conservative or operative), not on an intention to operate. My feeling is that waiting over a year may indeed be associated with a worse outcome, but unless the dropout rate of patients improving spontaneously during the waiting time is known, it cannot be proven.

doi:10.1302/0301-620X.87B6.16677

M. F. BROWN, PhD, FRCSEd(Orth)
University Hospital of North Staffordshire
Stoke-on-Trent, UK.

1. Ng LCL, Sell P. Predictive value of the duration of sciatica for lumbar discectomy. *J Bone Joint Surg [Br]* 2004;86-B:546-9.

Author's reply:

Sir,

We thank Mr Brown for his interest in our paper. In answer to his comment, we would consider it unlikely that readers of the Journal would draw an erroneous conclusion on the data presented. The title of the paper makes it clear that the cohort in question is

an operative one. Our article contributes to the literature regarding the optimum timing for operation in radicular pain.

The perfect answer to a question often needs to be balanced against the practical realities of clinical practice. The dropout rate to follow-up of a prospective cohort of non-operatively treated patients would be so high that the suggested comparison based on intention to treat could not be achieved in our clinics. A better level of evidence would be a randomised, controlled trial.

Readers of our paper will have noted that patients still improve in terms of pain after a year of symptoms, but that decline in disability as indicated by the outcome measures used showed less change with time, perhaps suggestive of the development of bio-psycho-social factors with chronic symptom duration. It is the authors' feeling, that this is the major factor, but it remains to be proven.

doi:10.1302/0301-620X.87B6.16678

P. SELL, BM, MSc, FRCS
L. NG, BM, MRCS
Leicester General Hospital,
Leicester, UK.

Internal fixation of complex fractures of the proximal humerus

Sir,

We read with interest the paper in the August 2004 issue entitled "Internal fixation of complex fractures of the proximal humerus" by Gerber, Werner and Vienne¹ and would like to raise the following points.

They describe the treatment of 34 shoulder fractures. There were nine different types of fracture and at least ten types of operation, undertaken by different surgeons with varying post-operative physiotherapy regimens. In some cases the same type of fracture pattern was treated with different operations. With so many variables we feel it is difficult to draw any valid conclusions from the results.

The aim of operative treatment was to achieve reduction of the fracture, and therefore a better functional outcome. We note that the four malunited fractures achieved a mean subjective shoulder value score of 88.75%. Those fractures without malunion had a mean subjective shoulder value score of 87.1%. The role of accurate reduction, therefore, seems still to be unproven with regard to function.

We would welcome the author's comments on these points.

doi:10.1302/0301-620X.87B6.16573

J. GIBBS
D. RICKETTS
The Princess Royal Hospital,
Haywards Heath, UK.

1. Gerber C, Werner CML, Vienne P. Internal fixation of complex fractures of the proximal humerus. *J Bone Joint Surg [Br]* 2004;86-B:848-55.