

van den Bekerom MP, Hilverdink EF, Sierevelt IN, et al. A comparison of hemiarthroplasty with total hip replacement for displaced intracapsular fracture of the femoral neck: a randomised controlled multicentre trial in patients aged 70 years and over.

J Bone Joint Surg [Br] 2010;92-B:1422-8.

The authors regret that Table II was published with some inaccurate values. The correct table is published below:

Table II. Operative details of all patients

| | HA* (n = 137) | THR† (n = 115) |
|---|-----------------|-----------------|
| Mean interval trauma to surgery in days (range) | 1.0 (0 to 10) | 1.0 (0 to 9) |
| Surgeon (consultant/resident) (%)‡ | 27 (25)/82 (75) | 51 (57)/38 (43) |
| Approach (%) | | |
| Posterolateral | 5 (4) | 22 (19) |
| Anterolateral | 132 (96) | 93 (81) |
| Anaesthesia (%) | | |
| Spinal | 92 (67) | 71 (62) |
| Epidural | 5 (4) | 11 (10) |
| Complete | 25 (18) | 30 (26) |
| Psoas block | 2 (11) | 0 (0) |
| Unknown | 13 (10) | 3 (2) |
| Blood loss (%) | | |
| < 500 ml | 111 (81) | 70 (61) |
| > 500 ml | 8 (6) | 25 (22) |
| Unknown | 18 (13) | 20 (17) |
| Duration of operation (%) | | |
| < 1 hour | 44 (32) | 10 (9) |
| 1 to 1.5 hours | 66 (48) | 65 (56) |
| > 1.5 hours | 15 (11) | 30 (26) |
| Unknown | 12 (9) | 10 (9) |
| Duration of hospital stay (range) | | |
| Mortality during hospital stay (%) | 17.1 (2 to 89) | 18.4 (4 to 86) |
| | 7 (5) | 5 (4) |

* HA, hemiarthroplasty

† THR, total hip replacement

‡ there were 26 unknown