

all biomechanical parameters, which was statistically significant for mean peak torque and total work of maximal repetition. In terms of flexion biomechanics, the differences were less striking, but the footprint group did have a statistically significant advantage in terms of work in the last-third of the repetitive testing period. However, clinical outcomes were good or excellent for patients in both groups. The groups are obviously relatively small and the procedures were performed in a single centre after a change in practice, i.e. the Endobutton procedures were performed first before the standard technique was changed to a footprint reconstruction. However, overall the authors feel the evidence strong enough to recommend adopting a footprint technique as standard.

## REFERENCES

1. Rao AJ, Chalmers PN, Cvetanovich GL, et al. Preoperative Doxycycline Does Not Reduce Propionibacterium acnes in Shoulder Arthroplasty. *J Bone Joint Surg Am.* 2018;100(11):958-964.
2. Chalmers PN, Beck L, Stertz I, Tashjian RZ. Hydrogen peroxide skin preparation reduces Cutibacterium acnes in shoulder

arthroplasty: a prospective, blinded, controlled trial. *J Shoulder Elbow Surg.* 2019;S1058-2746(19)30254-X.

3. Matsen FA 3rd, Whitson A, Neradilek MB, Pottinger PS, Bertelsen A, Hsu JE. Factors predictive of Cutibacterium periprosthetic shoulder infections: a retrospective study of 342 prosthetic revisions. *J Shoulder Elbow Surg.* 2019;S1058-2746(19)30548-8.
4. Robinson CM, Goudie EB, Murray IR, et al. Open reduction and plate fixation versus nonoperative treatment for displaced midshaft clavicular fractures: a multicenter, randomized, controlled trial. *J Bone Joint Surg Am.* 2013;95(17):1576-1584.
5. Ahrens PM, Garlick NI, Barber J, Tims EM; Clavicle Trial Collaborative Group. The Clavicle Trial: a multicenter randomized controlled trial comparing operative with nonoperative treatment of displaced midshaft clavicle fractures. *J Bone Joint Surg Am.* 2017;99(16):1345-1354.
6. Nicholson JA, Clement ND, Clelland AD, MacDonald D, Simpson AHRW, Robinson CM. Displaced midshaft clavicle fracture union can be accurately predicted with a delayed assessment at 6 weeks following injury: a prospective cohort study. *J Bone Joint Surg Am.* 2020;102(7):557-566.
7. Sarda P, Qaddori A, Nauschutz F, Boulton L, Nanda R, Bayliss N. Distal biceps tendon rupture: current concepts. *Injury.* 2013;44(4):417-420.
8. Ford SE, Andersen JS, Macknet DM, Connor PM, Loeffler BJ, Gaston RG. Major complications after distal biceps tendon

repairs: retrospective cohort analysis of 970 cases. *J Shoulder Elbow Surg.* 2018;27(10):1898-1906.

9. Rubinger L, Solow M, Johal H, Al-Asiri J. Return to work following a distal biceps repair: a systematic review of the literature. *J Shoulder Elbow Surg.* 2020;S1058-2746(19)30836-5.
10. Stull JD, Nicholson TA, Davis DE, Namdari S. Addition of 3% hydrogen peroxide to standard skin preparation reduces Cutibacterium acnes-positive culture rate in shoulder surgery: a prospective randomized controlled trial. *J Shoulder Elbow Surg.* 2020;29(2):212-216.
11. Ilyas AM, Mangan JJ, Graham J. Radial nerve palsy recovery with fractures of the humerus: an updated systematic review. *J Am Acad Orthop Surg.* 2020;28(6):e263-e269.
12. Liu YF, Hong CK, Hsu KL, et al. Intravenous administration of tranexamic acid significantly improved clarity of the visual field in arthroscopic shoulder surgery: a prospective, double-blind, and randomized controlled trial. *Arthroscopy.* 2020;36(3):640-647.
13. Ranebo MC, Björnsson Hallgren HC, Holmgren T, Adolfsson LE. Surgery and physiotherapy were both successful in the treatment of small, acute, traumatic rotator cuff tears: a prospective randomized trial. *J Shoulder Elbow Surg.* 2020;29(3):459-470.
14. Bellringer SF, Phadnis J, Human T, Redmond CL, Bain GI. Biomechanical comparison of transosseous cortical button and Footprint repair techniques for acute distal biceps tendon ruptures. *Shoulder Elbow.* 2020;12(1):54-62.

# Spine

**X-ref** For other Roundups in this issue that cross-reference with Spine see: Research Roundups 3 & 5.

## No benefit of early versus late ambulation after incidental durotomy in lumbar spine surgery

Incidental durotomy is common during elective spine surgery, with the incidence quoted as reaching up to 17% in some series. In the short-term, incidental durotomy can lead to prolonged hospital stay, meningitis, and low pressure symptoms. In terms of treatment, most will aim to repair the dural tears, which is typically followed by a period of prolonged mandatory bed rest (BR) to reduce the hydrostatic pressure at the repair site. Despite these precautions, persist leaks continue to plague these patients and can be a long-term bothersome problem. A group from **Zurich (Switzerland)** investigated the incidence of revision surgery following persist cerebrospinal fluid (CSF) leak after surgical inadvertent durotomy repair.<sup>1</sup> In

all, 60 patients who underwent lumbar spinal surgery and sustained an incidental durotomy repaired intraoperatively were randomized to receive early ambulation (EA) or 48 hours' BR postoperatively. There were no differences in age, sex, and preoperative Oswestry Disability Index. Two patients in each group required revision surgery due to persisting CSF leak, one in the BR group required antibiotic treatment for pneumonia, two were treated for postoperative lung oedema, and one sustained multiple embolic strokes. No medical complications were recorded in the EA group. Two patients in the BR group required surgery to treat wound complications. One patient in the EA group required surgery for a persistent wound leak without signs or symptoms of a CSF leak. The study concludes that there is limited benefit in prolonged BR after repair of an incidental durotomy, and that prolonged BR may be associated with significantly more medical complications. While there are clearly limitations to the design and sample size of this small randomized controlled trial, we applaud the authors for adding some high-quality evidence to this difficult debate.

## Is fusion needed for patients with lumbar spinal stenosis and substantial low back pain?

There are conflicting reports which explore the impact of lumbar decompression with or without fusion on lower back pain and leg pain when treating lumbar stenosis. As a result, it remains unclear whether patients with substantial back pain may improve with decompression alone, and whether the amount of postoperative leg pain affects the ultimate outcome of decompression surgery. This study performed in **Matsue-shi (Japan)** evaluates the effect of decompression alone for patients with severe lower back pain.<sup>2</sup> The team identified 222 patients who underwent spinal decompression without fusion, all of whom presented with a picture of low back pain and had completed one-year follow-up. The Numerical Rating Scale (NRS) and Japanese Orthopaedic Association Back Pain Evaluation Questionnaire (JOABPEQ) was completed by each patient both before surgery, and at three- and 12-months postoperatively. The overall mean NRS scores for lower back pain, leg pain, and numbness significantly improved at